

# 11 | Proof of van Kampen's Theorem

In the last chapter have seen the statement of van Kampen's theorem and some of examples of its applications. Our main goal in this chapter is to prove this result. For reference we state it here again:

**10.17 van Kampen Theorem.** *Let  $(X, x_0)$  be a pointed topological space and let  $U_1, U_2 \subseteq X$  be open sets such that  $X = U_1 \cup U_2$ . If the sets  $U_1$ ,  $U_2$ , and  $U_1 \cap U_2$  are path connected and  $x_0 \in U_1 \cap U_2$  then*

$$\pi_1(X, x_0) \cong \operatorname{colim}(\pi_1(U_1, x_0) \xleftarrow{i_{1*}} \pi_1(U_1 \cap U_2, x_0) \xrightarrow{i_{2*}} \pi_1(U_2, x_0))$$

where for  $k = 1, 2$  the homomorphism  $i_{k*}$  is induced by the inclusion map  $i_k: U_1 \cap U_2 \rightarrow U_k$ .

*Proof.* Here is some notation that will be useful.

- For simplicity we will denote  $U_1 \cap U_2$  by  $U_0$ .
- For  $k = 1, 2$  by  $i_k: U_0 \rightarrow U_k$  and  $j_k: U_k \rightarrow X$  we will denote the inclusion maps.
- If  $\omega$  is a loop in  $U_1$  then it represents an element of  $\pi_1(U_1, x_0)$  and an element of  $\pi_1(X, x_0)$ . In order to avoid such ambiguities we will write  $[\omega]_k$  to indicate an element of  $\pi_1(U_k, x_0)$  and  $[\omega]$  to indicate an element of  $\pi_1(X, x_0)$ .

The strategy of the proof will be as follows. Let  $P = \operatorname{colim}(\pi_1(U_1, x_0) \xleftarrow{i_{1*}} \pi_1(U_0, x_0) \xrightarrow{i_{2*}} \pi_1(U_2, x_0))$ . Recall that by Proposition 10.3  $P$  is the unique (up to an isomorphism) group that satisfies the following conditions:

- 1) for  $k = 1, 2$  there exists a homomorphism  $g_k: \pi_1(U_k, x_0) \rightarrow P$  such that  $g_1 i_{1*} = g_2 i_{2*}$ ;
- 2) for any group  $G$  and any homomorphisms  $h_k: \pi_1(U_k, x_0) \rightarrow G$  satisfying  $h_1 i_{1*} = h_2 i_{2*}$  there exists a unique homomorphism  $h: P \rightarrow G$  such that  $h g_k = h_k$  for  $k = 1, 2$ .

It follows that in order to prove van Kampen's theorem it will suffice to show that the group  $\pi_1(X, x_0)$  satisfies conditions 1) and 2). The first condition is satisfied by taking  $g_k = j_{k*}$  for  $k = 1, 2$ . In order to verify the second condition let  $h_1: \pi_1(U_1, x_0) \rightarrow G$  and  $h_2: \pi_1(U_2, x_0) \rightarrow G$  be homomorphisms satisfying  $h_1 i_{1*} = h_2 i_{2*}$ . We need to construct a homomorphism  $h: \pi_1(X, x_0) \rightarrow G$  such that the following diagram commutes:

$$\begin{array}{ccc}
 \pi_1(U_0, x_0) & \xrightarrow{i_{1*}} & \pi_1(U_1, x_0) \\
 \downarrow i_{2*} & & \downarrow j_{1*} \\
 \pi_1(U_2, x_0) & \xrightarrow{j_{2*}} & \pi_1(X, x_0) \\
 & \searrow h & \downarrow h_1 \\
 & & G
 \end{array}
 \quad (*)$$


$h_2$  (curved arrow from  $\pi_1(U_2, x_0)$  to  $G$ )

Moreover, we need to show that there is only one such homomorphism  $h$ .

The construction of  $h$  will use the following setup. For each point  $x \in X$  choose a path  $\sigma_x$  such that

- $\sigma_x(0) = x$  and  $\sigma_x(1) = x_0$ ;
- if  $x \in U_k$  for  $k \in \{0, 1, 2\}$  then  $\sigma_x$  is contained in  $U_k$ ;
- $\sigma_{x_0}$  is the constant path.

Such paths exist since by assumption  $U_0, U_1, U_2$  are path connected sets. If  $\omega$  is a path in  $X$  then the concatenation  $\bar{\sigma}_{\omega(0)} * \omega * \sigma_{\omega(1)}$  is a loop based at  $x_0$ :



We will denote this loop by  $\omega^\circ$  and call it the *loop completion* of  $\omega$ . Loop completion has the following properties:


- (i) If  $\omega$  is a path in  $U_k$  for  $k \in \{0, 1, 2\}$  then  $\omega^\circ$  is a loop in  $U_k$ . This holds since in such case  $\sigma_{\omega(0)}$  and  $\sigma_{\omega(1)}$  are paths in  $U_k$ .
- (ii) For any path  $\omega$  we have  $(\bar{\omega})^\circ \simeq \bar{\omega}^\circ$ .
- (iii) If  $\omega, \tau$  have the same endpoints and  $\omega \simeq \tau$  then  $\omega^\circ \simeq \tau^\circ$ , and so in  $\pi_1(X, x_0)$  we have  $[\omega^\circ] = [\tau^\circ]$ . Moreover, if  $\omega, \tau$  are paths in  $U_k$  and the homotopy between them is contained in  $U_k$  then in  $\pi_1(U_k, x_0)$  we have  $[\omega^\circ]_k = [\tau^\circ]_k$ .
- (iv) If  $\omega, \tau$  are paths such that  $\omega(1) = \tau(0)$  then  $(\omega * \tau)^\circ \simeq \omega^\circ * \tau^\circ$ . Indeed, we have

$$\omega^\circ * \tau^\circ = (\bar{\sigma}_{\omega(0)} * \omega * \sigma_{\omega(1)}) * (\bar{\sigma}_{\omega(1)} * \tau * \sigma_{\tau(1)}) \simeq \bar{\sigma}_{\omega(0)} * \omega * \tau * \sigma_{\tau(1)} \simeq (\omega * \tau)^\circ$$

Notice also that if both  $\omega$  and  $\tau$  are paths in  $U_k$  then the above homotopy of loops is contained in  $U_k$ . This means that  $[\omega^\circ]_k \cdot [\tau^\circ]_k = [(\omega * \tau)^\circ]_k$  in  $\pi_1(U_k, x_0)$ .

- (v) If  $\omega$  is a loop based at  $x_0$  then  $\omega \simeq \omega^\circ$ . To see this notice that in this case  $\sigma_{\omega(0)} = \sigma_{\omega(1)} = \sigma_{x_0}$ , and by assumption  $\sigma_{x_0}$  is the constant path. Therefore we have  $\omega^\circ = \bar{\sigma}_{x_0} * \omega * \sigma_{x_0} \simeq \omega$ .


We are now ready to describe the construction of the homomorphism  $h$  in the diagram (\*). Let  $\omega: [0, 1] \rightarrow X$  be a loop. The sets  $\omega^{-1}(U_1)$  and  $\omega^{-1}(U_2)$  form an open cover of the interval  $[0, 1]$ . Using the Lebesgue number of this cover we obtain that there exists an  $n$ -tuple of numbers  $(s_0, \dots, s_n)$  such that  $0 = s_0 < s_1 < \dots < s_n = 1$  and  $\omega([s_{i-1}, s_i])$  is contained in either  $U_1$  or  $U_2$  for each  $i = 1, \dots, n$ . Let  $\omega_i: [0, 1] \rightarrow X$  be the path given by  $\omega_i(s) = \omega(s_{i-1}s + s_i(1 - s))$ . This path coincides with the restriction of  $\omega$  to the subinterval  $[s_{i-1}, s_i]$ :



We have  $\omega \simeq \omega_1 * \omega_2 * \dots * \omega_n$ . By the properties of loop completion we get:

$$\omega \simeq \omega^\circ \simeq \omega_1^\circ * \omega_2^\circ * \dots * \omega_n^\circ$$

Moreover, if  $\omega_i$  is contained in  $U_1$  then  $\omega_i^\circ$  is a loop in  $U_1$ , and likewise for  $U_2$ .



Let  $k(i)$  denote either 1 or 2 depending if  $\omega_i^\circ$  is a loop in  $U_1$  or  $U_2$ . Define an element  $\tilde{h}(\omega) \in G$  by

$$\tilde{h}(\omega) = h_{k(1)}([\omega_1^\circ]_{k(1)}) \cdot h_{k(2)}([\omega_2^\circ]_{k(2)}) \cdot \dots \cdot h_{k(n)}([\omega_n^\circ]_{k(n)}) \quad (**)$$

There are two ambiguities in this formula. First, if  $\omega_i$  is a path entirely contained in  $U_0$  then  $\omega_i^\circ$  is a loop in  $U_0$  and in such case we can take  $k(i)$  to be either 1 or 2. This however does not matter since in such case we have  $[\omega_i]_1 = i_{1*}([\omega_i]_0)$  and  $[\omega_i]_2 = i_{2*}([\omega_i]_0)$ , and since  $h_1 i_{1*} = h_2 i_{2*}$  we get

$$h_1([\omega_i^\circ]_1) = h_1 i_{1*}([\omega_i^\circ]_0) = h_2 i_{2*}([\omega_i^\circ]_0) = h_2([\omega_i^\circ]_1)$$

The second ambiguity comes from the fact that the formula (\*\*) uses subdivision of  $\omega$  into paths  $\omega_i$ , and the value of  $\tilde{h}(\omega)$  could change if we change the subdivision. To see that this is not the case consider the subdivision  $\omega \simeq \omega_1 * \dots * \omega_n$  that comes from an  $n$ -tuple of numbers  $\underline{s} = (s_0, \dots, s_n)$ , and let  $s_+$  be a number such that  $s_{i-1} < s_+ < s_i$  for some  $1 \leq i \leq n$ . The  $(n+1)$ -tuple  $\underline{s}' = (s_1, \dots, s_{i-1}, s_+, s_{i+1}, \dots, s_n)$  produces the subdivision

$$\omega \simeq \omega_1 * \dots * \omega_{i-1} * \tau_1 * \tau_2 * \omega_{i+1} * \dots * \omega_n$$

where  $\tau_1 * \tau_2 \simeq \omega_i$ . By the properties of loop completion  $[\omega^\circ]_{k(i)} = [\tau_1^\circ]_{k(i)} \cdot [\tau_2^\circ]_{k(i)}$ , and thus we get

$$h_{k(i)}([\omega_i^\circ]_{k(i)}) = h_{k(i)}([\tau_1^\circ]_{k(i)} \cdot [\tau_2^\circ]_{k(i)}) = h_{k(i)}([\tau_1^\circ]_{k(i)}) \cdot h_{k(i)}([\tau_2^\circ]_{k(i)})$$

This shows the the value of  $\tilde{h}(\omega)$  does not depend on whether we compute it using  $\underline{s}$  or  $\underline{s}'$ . Arguing inductively we obtain that if an  $m$ -tuple  $\underline{s}'$  is obtained by adding some number of elements to an  $n$ -tuple  $\underline{s}$  then the value of  $\tilde{h}(\omega)$  computed using  $\underline{s}'$  is the same as the value computed using  $\underline{s}$ . Finally, given an arbitrary  $n$ -tuple  $\underline{s}$  and an  $m$ -tuple  $\underline{s}'$  that produce two subdivisions of  $\omega$  we can find an  $r$ -tuple  $\underline{s}''$  that can be obtained from each of  $\underline{s}$  and  $\underline{s}'$  by inserting some additional elements. The argument above shows then that the values of  $\tilde{h}(\omega)$  computed using  $\underline{s}$  and  $\underline{s}'$  must be equal since they are both equal to the value computed using  $\underline{s}''$ .

Our goal will be to show that  $\tilde{h}(\omega)$  depends only on the homotopy class of  $\omega$ :

*Claim.* If  $\omega, \tau$  are loops in  $X$  and  $\omega \simeq \tau$  then  $\tilde{h}(\omega) = \tilde{h}(\tau)$ .

Assuming for a moment this claim holds notice that it allows us to define a function  $h: \pi_1(X, x_0) \rightarrow G$  by  $h([\omega]) = \tilde{h}(\omega)$ . Notice also that this function has the following properties:

- 1)  $h$  is a homomorphism;
- 2)  $h$  makes the diagram (\*) into a commutative diagram;
- 3)  $h$  is the only homomorphism that makes the diagram (\*) commute.

Indeed, to see 1) observe that if  $\omega$  and  $\omega'$  are two loops in  $X$  with subdivisions  $\omega \simeq \omega_1 * \dots * \omega_n$  and  $\omega' \simeq \omega'_1 * \dots * \omega'_m$  then  $\omega * \omega'$  has a subdivision

$$\omega * \omega' \simeq \omega_1 * \dots * \omega_n * \omega'_1 * \dots * \omega'_m$$

Using this observation and the definition of  $h$  we get that  $h([\omega] \cdot [\omega']) = h([\omega]) \cdot h([\omega'])$ .

To verify 2) notice that if  $\omega$  is a loop in  $U_k$  for either  $k = 1$  or  $k = 2$  then we don't need to subdivide it, i.e. we can take  $\omega = \omega_1$ . Since  $\omega \simeq \omega^\circ$  the formula (\*\*) gives


$$h([\omega]) = \tilde{h}(\omega) = h_k([\omega]_k)$$

and since  $[\omega] = j_{k*}([\omega]_k)$  we obtain  $h j_{k*} = h_k$ .

Finally, to see 3) notice that we have shown that any element  $[\omega] \in \pi_1(X, x_0)$  can be written as a product  $[\omega] = [\omega_1^\circ] \cdot \dots \cdot [\omega_n^\circ]$  where  $\omega_i$  is a loop in  $U_{k(i)}$  for either  $k(i) = 1$  or  $k(i) = 2$ . For each

$i = 1, \dots, n$  we have  $[\omega_i^\circ] = j_{k(i)*}([\omega_i]_{k(i)})$ . Thus in order to get the identity  $h j_{k(i)*} = h_{k(i)}$  we must set  $h([\omega_i^\circ]) = h_{k(i)}([\omega_i]_{k(i)})$ . Since homomorphisms preserve products we are forced to define  $h([\omega])$  by the formula (\*\*).

The above observations show that once we verify that  $\tilde{h}$  is a homotopy invariant function the proof of van Kampen's Theorem will be complete. Assume then that  $\omega, \tau$  are loops in  $X$  and that there exists a homotopy  $h: [0, 1] \times [0, 1] \rightarrow X$  between them:  $h_0 = \omega, h_1 = \tau$ . We will consider first a special case, and assume in addition that there exists numbers  $0 = s_0 < s_1 < \dots < s_n = 1$  such that the homotopy  $h$  maps each rectangle  $[s_{i-1}, s_i] \times [0, 1]$  either into  $U_1$  or into  $U_2$ . In particular this gives a subdivisions  $\omega \simeq \omega_1 * \dots * \omega_n$  and  $\tau \simeq \tau_1 * \dots * \tau_n$  where  $\omega_i$  and  $\tau_i$  are restrictions of  $\omega$  and  $\tau$  to the subinterval  $[s_{i-1}, s_i]$ . For  $i = 0, \dots, n$  denote by  $\gamma_i$  the path given by  $\gamma(t) = h(s_i, t)$ . Notice that  $\gamma_0$  and  $\gamma_n$  are constant paths at  $x_0$ .



Let  $k(i)$  denote either 1 or 2 depending if  $h([s_{i-1}, s_i] \times [0, 1])$  is contained in  $U_1$  or  $U_2$ . Notice that for each  $i$  the paths  $\omega_i, \tau_i, \gamma_{i-1}$ , and  $\gamma_i$  are paths in  $U_{k(i)}$ . Moreover, using the restriction of  $h$  to  $[s_{i-1}, s_i] \times [0, 1]$  we obtain that the path  $\omega_i$  is homotopic to  $\gamma_{i-1} * \tau_i * \bar{\gamma}_i$  via a homotopy contained in  $U_{k(i)}$ . Using the properties of loop completion we obtain

$$[\omega_i^\circ]_{k(i)} = [(\gamma_{i-1} * \tau_i * \bar{\gamma}_i)^\circ]_{k(i)} = [\gamma_{i-1}^\circ]_{k(i)} \cdot [\tau_i^\circ]_{k(i)} \cdot [\gamma_i^\circ]_{k(i)}^{-1}$$

This identity and the formula (\*\*) gives:

$$\begin{aligned} \tilde{h}(\omega) &= h_{k(1)}([\omega_1^\circ]_{k(1)}) \cdot h_{k(2)}([\omega_2^\circ]_{k(2)}) \cdot \dots \cdot h_{k(n)}([\omega_n^\circ]_{k(n)}) \\ &= h_{k(1)}([\gamma_0^\circ]_{k(1)}) \cdot h_{k(1)}([\tau_1^\circ]_{k(1)}) \cdot h_{k(1)}([\gamma_1^\circ]_{k(1)})^{-1} \\ &\quad \cdot h_{k(2)}([\gamma_1^\circ]_{k(2)}) \cdot h_{k(2)}([\tau_2^\circ]_{k(2)}) \cdot h_{k(2)}([\gamma_2^\circ]_{k(2)})^{-1} \cdot \dots \\ &\quad \cdot h_{k(n)}([\gamma_{n-1}^\circ]_{k(n)}) \cdot h_{k(n)}([\tau_n^\circ]_{k(n)}) \cdot h_{k(n)}([\gamma_n^\circ]_{k(n)})^{-1} \end{aligned} \quad (***)$$


Notice that  $h_{k(1)}([\gamma_1^\circ]_{k(1)})$  and  $h_{k(n)}([\gamma_n^\circ]_{k(n)})$  are trivial elements of  $G$  since  $\gamma_1^\circ$  and  $\gamma_n^\circ$  are constant loops. Also, for  $i = 1, \dots, n-1$  we have  $h_{k(i)}([\gamma_i^\circ]_{k(i)}) = h_{k(i+1)}([\gamma_i^\circ]_{k(i+1)})$ . This is obvious if  $k(i) = k(i+1)$ . If  $k(i) \neq k(i+1)$  then this identity still holds since in such case  $\gamma_i$  (and thus also  $\gamma_i^\circ$ ) is contained in  $U_0$  and then

$$h_1([\gamma_i^\circ]_1) = h_1 i_{1*}([\gamma_i^\circ]_0) = h_2 i_{2*}([\gamma_i^\circ]_0) = h_2([\gamma_i^\circ]_2)$$

Using these observations we can simplify the equation (\*\*) to

$$\tilde{h}(\omega) = h_{k(1)}([\tau_1^\circ]_{k(1)}) \cdot h_{k(2)}([\tau_2^\circ]_{k(2)}) \cdot \dots \cdot h_{k(n)}([\tau_n^\circ]_{k(n)}) = \tilde{h}(\tau)$$

It remains to consider the general case when we are given two loops  $\omega$  and  $\tau$  in  $X$  such that  $\omega \simeq \tau$ . Let  $h: [0, 1] \times [0, 1] \rightarrow X$  be a homotopy with  $h_0 = \omega$  and  $h_1 = \tau$ . The sets  $h^{-1}(U_1)$  and  $h^{-1}(U_2)$  form an open cover of  $X$ . Using the Lebesgue number of this cover we can find numbers  $0 = s_0 < s_1 < \dots < s_n = 1$  and  $0 = t_0 < t_1 < \dots < t_m = 1$  such that  $h$  maps each rectangle  $[s_{i-1}, s_i] \times [t_{j-1}, t_j]$  either into  $U_1$  or into  $U_2$ . Let  $\delta_i$  be the path given by  $\delta_i(s) = h(s, t_i)$ :



Notice that the restriction of  $h$  to the rectangle  $[0, 1] \times [t_{i-1}, t_i]$  gives a homotopy between  $\delta_{i-1}$  and  $\delta_i$ , and moreover this homotopy is of the form considered in the special case above. Therefore for each  $i = 1, \dots, m$  we have  $\tilde{h}(\delta_{i-1}) = \tilde{h}(\delta_i)$ . As a consequence we obtain

$$\tilde{h}(\omega) = \tilde{h}(\delta_0) = \tilde{h}(\delta_1) = \dots = \tilde{h}(\delta_m) = \tilde{h}(\tau)$$

□

Theorem 10.17 can be generalized to the case where the space  $X$  is covered by more than two (and possibly infinitely many) open sets:

**11.1 Theorem.** Let  $(X, x_0)$  be a pointed topological space and let  $\{U_i\}_{i \in I}$  be an open cover of  $X$  such that  $x_0 \in U_i$  for all  $i \in I$ . For  $i, j \in I$  let  $f_{ij}: U_i \cap U_j \rightarrow U_i$  denote the inclusion map. If the set  $U_i \cap U_j \cap U_k$  is path connected for all  $i, j, k \in I$  then


$$\pi_1(X, x_0) \cong *_{i \in I} \pi_1(U_i, x_0) / N$$

where  $N$  is the normal subgroup of  $*_{i \in I} \pi_1(U_i, x_0)$  generated by all elements of the form  $f_{ij}([\omega]) \cdot f_{ji}([\omega])^{-1}$  for  $i, j \in I$  and  $[\omega] \in \pi_1(U_i \cap U_j, x_0)$ .

Here is one application of the generalized van Kampen Theorem:

**11.2 Proposition.** Let  $\{(X_i, x_i)\}_{i \in I}$  be a family of path connected pointed spaces, and let  $\bigvee_{i \in I} X_i$  be the space obtained by identifying the basepoints  $x_i \sim x_j$  for all  $i, j \in I$ . Assume that for each  $i \in I$  there exists a set  $V_i \subseteq X_i$  such that  $V_i$  is open in  $X_i$ ,  $x_i \in V_i$ , and the one-point space  $\{x_i\}$  is a deformation retract of  $V_i$ . Then

$$\pi_1(\bigvee_i X_i) \cong *_{i \in I} \pi_1(X_i, x_i)$$



*Proof.* We will denote by  $\bar{x}$  the point of  $\bigvee_{i \in I} X_i$  obtained by identifying the points  $x_i \in X_i$ . For  $i \in I$  let  $U_i = X_i \vee \bigvee_{i \in I} V_i$ . The sets  $U_i$  are path connected and they form an open cover  $\bigvee_i X_i$ . Moreover for  $j, k, l \in I$  we have

$$U_j \cap U_k \cap U_l = \begin{cases} U_j & \text{if } j = k = l \\ \bigvee_{i \in I} V_i & \text{otherwise} \end{cases}$$

so  $U_j \cap U_k \cap U_l$  is path connected for all  $j, k, l$ . Using Theorem 11.1 we obtain

$$\pi_1(\bigvee_{i \in I} X_i, \bar{x}) \cong *_{i \in I} \pi_1(U_i, x_i) / N$$

where  $N$  is the normal group generated by elements  $f_{ij*}([\omega]) \cdot f_{ji*}([\omega])^{-1}$  for  $[\omega] \in \pi_1(U_i \cap U_j, \bar{x})$ . As in Theorem 11.1 by  $f_{ij}$  we denote here the inclusion map  $f_{ij}: U_i \cap U_j \rightarrow U_{ij}$ . For  $i \in I$  let  $h_i: V_i \times [0, 1] \rightarrow V_i$  the deformation retraction of  $V_i$  onto  $\{x_i\}$ . These maps define a deformation retraction of the space  $\bigvee_{i \in I} V_i$  onto  $\{\bar{x}\}$ . This shows that the space  $\bigvee_{i \in I} V_i$  is contractible and so  $\pi_1(\bigvee_{i \in I} V_i, \bar{x}) = \{1\}$ . Since for all  $i, j \in I$  such that  $i \neq j$  we have  $U_i \cap U_j = \bigvee_{i \in I} V_i$  it follows that the group  $N$  is trivial and so  $\pi_1(\bigvee_{i \in I} X_i, \bar{x}) \cong *_{i \in I} \pi_1(U_i, x_i)$ . Finally, using again the deformation retractions  $h_i$  we can construct for each  $j \in I$  a deformation retraction of  $U_j$  onto  $X_j$ . This shows that  $\pi_1(U_j, \bar{x}) \cong \pi_1(X_j, \bar{x})$ , and so we get

$$\pi_1(\bigvee_{i \in I} X_i, \bar{x}) \cong *_{i \in I} \pi_1(U_i, \bar{x}) \cong *_{i \in I} \pi_1(X_i, \bar{x})$$

□

**11.3 Example.** Let  $s_0$  be a basepoint of  $S^1$ . Taking  $V = S^1 \setminus \{s_1\}$  where  $s_1 \in S^1$  is a point different from  $s_0$  we obtain an open neighborhood of  $s_0$  which deformation retracts onto  $\{s_0\}$ . Thus using Proposition 11.2 we get

$$\pi_1(\bigvee_{i \in I} S^1) \cong *_{i \in I} \pi_1(S^1) \cong *_{i \in I} \mathbb{Z}$$

## Exercises to Chapter 11